

Top FB asymmetry at the Tevatron and possible new physics signatures

Pyungwon Ko

Korea Institute for Advanced Study (KIAS)

in collaboration with
Dr. J.-S. Lee, Dr. D.W. Jung (and Dr. S.H. Nam)

based on arXiv:0912.1105, PLB 2010.06.040 and
arXiv:1011.5976 and work in preparation

< KEK-PH 2011, Mar. 2-5 (2011) >


Outline

- 1 Introduction
- 2 Effective Lagrangian Approach
 - Forward-Backward Asymmetry
 - Helicity Amplitude
 - Spin-Spin Correlation
- 3 Models including new resonances explicitly
 - Spin-1 Resonances
 - Spin-0 Resonances
 - Wilson Coefficients from Resonances
 - Examples of Resonances
- 4 Longitudinal Pol of (anti)top quark
 - P violation and Longitudinal Pol's
- 5 Summary


Introduction

Top pair production via strong interaction

- Tevatron:



- LHC:



Introduction

Motivation

- Top physics has begun to enter a new era after its first discovery, due to the high luminosity achieved at the Tevatron, and precision study will be possible at the LHC in the coming years.
- Forward-backward asymmetry A_{FB}^t in $t\bar{t}$ production has been off the SM prediction ($\sim 0.078(9)$) by 2σ in the $t\bar{t}$ rest frame (CDF2008):

$$A_{\text{FB}}^t \equiv \frac{N_t(\cos\theta \geq 0) - N_{\bar{t}}(\cos\theta \geq 0)}{N_t(\cos\theta \geq 0) + N_{\bar{t}}(\cos\theta \geq 0)} = 0.24 \pm 0.13 \pm 0.04$$

- This $\sim 2\sigma$ deviation stimulated some speculations on new physics scenarios
- We adopt a model independent approach using effective Lagrangian in order to accommodate the current measurement of A_{FB}^t , since there is no clear evidence for any new particles coupling to top at the Tevatron

Introduction

- New CDF data with 5.6 fb^{-1} presented this summer

$$0.158 \pm 0.072 \pm 0.017$$


- Less deviation than before
→ Any new physics scale is probably too high to be explored directly at the Tevatron
- Still interesting to speculate what type of new physics can modify top physics at what level
- In fact, our approach based on the effective lagrangian could be more useful in this case, than other approaches
- Also could be used to set **substructure scale of top quark**, as in the light quark system

Introduction

SM contribution to A_{FB}

Series of papers by Kühn and Rodrigo

- LO: top quark production angle is symmetric with respect to beam direction.
- NLO: asymmetry due to interference effects.



Introduction


Related works as of Summer 2010 (Not updated afterwards)

- SM predictions: Kühn and Rodrigo
 - ▶ Interference between tree (one gluon exchange) and one-loop (two gluon exchange)
 - ▶ (anti)quark-gluon scattering into $t\bar{t}g$
 - ▶ initial (final) gluon emission $q\bar{q} \rightarrow t\bar{t}g$
- Axigluon: Godbole et al.; Rodrigo et al.; Frampton, Shu, Wang; Chivukular, Simmons, CP Yuan
- Extra Z' : Jung, Murayama, Pierce, Wells
- Extra W' : Cheung, Keung, TC Yuan
- RS KK gluon: Djouadi et al.
- Color sextet or antitriplet: Tait et al.; CH Chen et al.; Berger et al.;
- RPV and LR model: Cao, Heng, Wu, Yang
- Comprehensive study: Cao, McKeen, Rosner, Shaughnessy, Wagner
- Effective Lagrangian Approach : this talk
- Apologies to those who are not listed

Introduction

Wisdom from EW sector

- The first evidence of asymmetry was found in angular distribution of muons from e^+e^- collisions at PETRA in the 80's ($\sqrt{s} \sim 30$ GeV, well below the Z^0 pole)



- Source of A_{FB} is a term linear in $\cos \theta$ from interference between γ or Z vector coupling and the axial vector Z coupling.

Effective Lagrangian Approach

Dim-6 Contact Interaction

- $t\bar{t}$ production at the Tevatron dominated by $q\bar{q}$ channel
- Enough to consider dimension-6 four-quark operators **assuming new physics scale is high enough**:

$$\mathcal{L}_6 = \frac{g_s^2}{\Lambda^2} \sum_{A,B} \left[C_{1q}^{AB} (\bar{q}_A \gamma_\mu q_A) (\bar{t}_B \gamma^\mu t_B) + C_{8q}^{AB} (\bar{q}_A T^a \gamma_\mu q_A) (\bar{t}_B T^a \gamma^\mu t_B) \right]$$

where

$$T^a = \lambda^a/2, \quad \{A, B\} = \{L, R\}, \quad L, R \equiv (1 \mp \gamma_5)/2 \quad (q = u, d, s, c, b)$$

- Other d=6 operators are all reducible to the above operators after Fierzing (Hill and Parke 1994)
- We ignore flavor changing dim-6 operators such as $\bar{d}_R \gamma^\mu s_R \bar{t}_R \gamma_\mu t_R$, since those contributions to the $t\bar{t}$ production cross section will be of a order $1/\Lambda^4$

Effective Lagrangian Approach

Digress on Top Substructure

- Our approach is useful even if the A_{FB} approaches the SM prediction
- This contact term used to explore light quark substructures
- Similar analysis has been done for light quark and lepton systems using $\bar{q}q\bar{q}q$, $\bar{q}q\bar{l}l$, and $\bar{l}l\bar{l}l$, with various Dirac and color structures
- One can do exactly the same analysis for top compositeness scale
- The scale Λ in our effective lagrangian could be interpreted as **Compositeness scale of a top quark seen by a light quark**
- Bound on Λ can be derived from $M_{t\bar{t}}$ or p_T^t distributions at the Tevatron

Effective Lagrangian Approach

Helicity Amplitude Squared

- The squared helicity amplitude is given by

$$\begin{aligned} |\overline{\mathcal{M}(t_L \bar{t}_L + t_R \bar{t}_R)}|^2 &= \frac{4 g_s^4}{9 \hat{s}} m_t^2 \left[2 + \frac{\hat{s}}{\Lambda^2} (C_1 + C_2) \right] s_{\hat{\theta}}^2 \\ |\overline{\mathcal{M}t_L \bar{t}_R + t_R \bar{t}_L}|^2 &= \frac{2 g_s^4}{9} \left[\left(1 + \frac{\hat{s}}{2\Lambda^2} (C_1 + C_2) \right) (1 + c_{\hat{\theta}}^2) \right. \\ &\quad \left. + \hat{\beta}_t \left(\frac{\hat{s}}{\Lambda^2} (C_1 - C_2) \right) c_{\hat{\theta}} \right] \end{aligned}$$


where

$$\begin{aligned} C_1 &\equiv C_{8q}^{LL} + C_{8q}^{RR}, & C_2 &\equiv C_{8q}^{LR} + C_{8q}^{RL} \\ \hat{\beta}_t^2 &= 1 - 4m_t^2/\hat{s}, & s_{\hat{\theta}} &\equiv \sin \hat{\theta}, & c_{\hat{\theta}} &\equiv \cos \hat{\theta} \end{aligned}$$

- The term linear in $\cos \hat{\theta}$ could generate the forward-backward asymmetry which is proportional to $\Delta C \equiv C_1 - C_2$.

Effective Lagrangian Approach

Allowed region in the (C_1, C_2) plane



Effective Lagrangian Approach

Validity of our approach

- Our Validity Criteria:
 - ▶ $\sigma_{\text{int}} < r\sigma_{\text{SM}}$ (straight line)
 - ▶ $\sigma_{\text{NP}} < r\sigma_{\text{int}}$ (ellipses passing through the origin)
 - ▶ $\sigma_{\text{NP}} < r^2\sigma_{\text{SM}}$ (ellipses centered at the origin)
- Take $r = 0.3$, $r = 0.5$, and $r = 1.0$
- Our predictions pass these validity criteria even for $r = 0.3$, and could be considered reliable
- Another Issue: Violation of Unitarity by dim-6 op's **Any nonrenormalizable interactions violate “unitarity”**, which is very subtle issue at hadron colliders, since $\sqrt{\hat{s}}$ is not fixed
- Our criteria is hopefully stronger than unitarity constraint

Effective Lagrangian Approach

Validity of our approach

σ_{NP} is obtained using


$$\begin{aligned} \overline{|\mathcal{M}_{\text{NP}}|^2} &= \frac{4g_s^4 \hat{s}^2}{9\hat{s}^2 4\Lambda^4} \\ &\times \left\{ [9 ((C_{1q}^{LL})^2 + (C_{1q}^{RR})^2) + 2 ((C_{8q}^{LL})^2 + (C_{8q}^{RR})^2)] (\hat{u} - m_t^2)^2 \right. \\ &\quad + [9 ((C_{1q}^{RL})^2 + (C_{1q}^{LR})^2) + 2 ((C_{8q}^{RL})^2 + (C_{8q}^{LR})^2)] (\hat{t} - m_t^2)^2 \\ &\quad \left. + [9 (C_{1q}^{LL} C_{1q}^{LR} + C_{1q}^{RR} C_{1q}^{RL}) + 2 (C_{8q}^{LL} C_{8q}^{LR} + C_{8q}^{RR} C_{8q}^{RL})] (2\hat{s}m_t^2) \right\}, \end{aligned}$$

where

$$\hat{u} - m_t^2 = -\hat{s}(1 + \hat{\beta}_t c_{\hat{\theta}})/2, \quad \hat{t} - m_t^2 = -\hat{s}(1 - \hat{\beta}_t c_{\hat{\theta}})/2$$


Effective Lagrangian Approach

Validity region




Effective Lagrangian Approach

Validity region with the updated data



Effective Lagrangian Approach

$M_{\bar{t}t}$ distribution



Effective Lagrangian Approach

Spin-Spin Correlation

- chiral structure of new physics affecting $q\bar{q} \rightarrow t\bar{t}$ is also sensitive to the top quark spin-spin correlation (in the helicity basis):

$$-K = C = \frac{\sigma(t_L\bar{t}_L + t_R\bar{t}_R) - \sigma(t_L\bar{t}_R + t_R\bar{t}_L)}{\sigma(t_L\bar{t}_L + t_R\bar{t}_R) + \sigma(t_L\bar{t}_R + t_R\bar{t}_L)}$$

- SM prediction for helicity basis: $K = 0.47$ (LO) and 0.352 (NLO) [Bernreuther et al., NPB (2004)]

- New CDF data:


$$K = 0.48 \pm 0.48 \pm 0.22$$

- New physics should have **chiral couplings both to light quarks and top quark** \longrightarrow **P must be broken**
- Any new observable ?

Longitudinal top quark polarization

Effective Lagrangian Approach

New Spin-Spin Correlation C_{FB}



Effective Lagrangian Approach

Proposing a “NEW” Spin-Spin Correlation

- The usual C is correlated with $\sigma_{t\bar{t}}$, and not to A_{FB}
- We propose a new spin-spin correlation C_{FB} : Separate the events in forward and backward directions, and form C_{FB}


$$C_{FB} \equiv C(\cos \theta \geq 0) - C(\cos \theta \leq 0)$$

$C(\cos \theta \geq 0(\leq 0))$ implies the cross sections in the numerator of C are obtained for the forward (backward) region: $\cos \theta \geq 0(\leq 0)$

- Advantages of the new C_{FB} :
 - ▶ Larger spin-spin correlation
 - ▶ Stronger correlation with A_{FB}
- This new C_{FB} could be also useful for testing the QCD in the top sector

Effective Lagrangian Approach

Spin-Spin Correlation



Spin-1 Resonances

- One can consider the following interactions of quarks with spin-1 flavor-conserving (changing) color-singlet $V_1(\tilde{V}_1)$ and color-octet $V_8^a(\tilde{V}_8^a)$ vectors ($A = L, R$) relevant to A_{FB}^t :

$$\begin{aligned}\mathcal{L}_V = & g_s V_1^\mu \sum_A \left[g_{1q}^A (\bar{q}_A \gamma_\mu q_A) + g_{1t}^A (\bar{t}_A \gamma_\mu t_A) \right] \\ & + g_s V_8^{a\mu} \sum_A \left[g_{8q}^A (\bar{q}_A \gamma_\mu T^a q_A) + g_{8t}^A (\bar{t}_A \gamma_\mu T^a t_A) \right] \\ & + g_s \left[\tilde{V}_1^\mu \sum_A \tilde{g}_{1q}^A (\bar{t}_A \gamma_\mu q_A) + \tilde{V}_8^{a\mu} \sum_A \tilde{g}_{8q}^A (\bar{t}_A \gamma_\mu T^a q_A) + \text{h.c.} \right]\end{aligned}$$

Spin-0 Resonances

- Following interactions of quarks with spin-0 flavor-changing color-singlet \tilde{S}_1 and color-octet \tilde{S}_8^a scalars could also contribute to A_{FB}^t :

$$\mathcal{L}_{\tilde{S}} = g_s \left[\tilde{S}_1 \sum_A \tilde{\eta}_{1q}^A (\bar{t} A q) + \tilde{S}_8^a \sum_A \tilde{\eta}_{8q}^A (\bar{t} A T^a q) + \text{h.c.} \right]$$

- One can also consider color-triplet S_k^γ and color-sextet scalars $S_{ij}^{\alpha\beta}$ with minimal flavor violating interactions with the SM quarks (Arnold, Pospelov, Trott, Wise):

$$\mathcal{L}_S = g_s \left[\frac{\eta_3}{2} \epsilon_{\alpha\beta\gamma} \epsilon^{ijk} u_{iR}^\alpha u_{jR}^\beta S_k^\gamma + \eta_6 u_{iR}^\alpha u_{jR}^\beta S_{ij}^{\alpha\beta} + \text{h.c.} \right]$$

Wilson Coefficients from Resonances

- After integrating out the heavy vectors and scalars, we obtain the Wilson coefficients as follows:

$$\begin{aligned}\frac{C_{8q}^{LL}}{\Lambda^2} &= -\frac{1}{m_V^2} g_{8q}^L g_{8t}^L - \frac{1}{m_{\tilde{V}}^2} \left[2|\tilde{g}_{1q}^L|^2 - \frac{1}{N_c} |\tilde{g}_{8q}^L|^2 \right] \\ \frac{C_{8q}^{RR}}{\Lambda^2} &= -\frac{1}{m_V^2} g_{8q}^R g_{8t}^R - \frac{1}{m_{\tilde{V}}^2} \left[2|\tilde{g}_{1q}^R|^2 - \frac{1}{N_c} |\tilde{g}_{8q}^R|^2 \right] - \frac{|\eta_3|^2}{m_{S_3}^2} + \frac{2|\eta_6|^2}{m_{S_6}^2} \\ \frac{C_{8q}^{LR}}{\Lambda^2} &= -\frac{1}{m_V^2} g_{8q}^L g_{8t}^R - \frac{1}{m_S^2} \left[|\tilde{\eta}_{1q}^L|^2 - \frac{1}{2N_c} |\tilde{\eta}_{8q}^L|^2 \right] \\ \frac{C_{8q}^{RL}}{\Lambda^2} &= -\frac{1}{m_V^2} g_{8q}^R g_{8t}^L - \frac{1}{m_S^2} \left[|\tilde{\eta}_{1q}^R|^2 - \frac{1}{2N_c} |\tilde{\eta}_{8q}^R|^2 \right]\end{aligned}$$

Examples of Resonances

- Axigluon model corresponding to flavor universal chiral couplings (Pati and Salam 1975): $g_{8q}^L = g_{8t}^L = -g_{8q}^R = -g_{8t}^R = 1$

- New gauge boson Z' with dominant coupling to $u - t$ (Jung, Murayama, Pierce, and Wells 2009):

$$V_1 = \tilde{V}_1 = Z', \quad g_s \tilde{g}_{1q}^R = g_X, \quad g_s g_{1q}^R = g_X \epsilon_U \quad (|\epsilon_U| \lesssim 1)$$

- New charged gauge boson W'^{\pm} contributions (Cheung, Keung, and Yuan 2009): $\tilde{V} = W', \quad g_s \tilde{g}_{1q}^A = g' g_A$

- Some RS scenarios with large flavor mixing in the right-handed quark sector (Aquino et al 2007; Agashe et al 2008):


$$g_{8q}^L = g_{8q}^R = g_{8b}^R \simeq -0.2, \quad g_{8t}^L = g_{8b}^L \simeq (1 \sim 2.8)$$

$$g_{8t}^R \simeq (1.5 \sim 5), \quad \tilde{g}_{8q}^L \simeq V_{tq}, \quad \tilde{g}_{8q}^R \simeq 1$$

Scores for each model

New particle	couplings	C_1	C_2	1σ favor
V_8 (spin-1 FC octet)	$g_{8q,8t}^{L,R}$	indefinite	indefinite	✓
\tilde{V}_1 (spin-1 FV singlet)	$\tilde{g}_{1q}^{L,R}$	–	0	×
\tilde{V}_8 (spin-1 FV octet)	$\tilde{g}_{8q}^{L,R}$	+	0	✓
\tilde{S}_1 (spin-0 FV singlet)	$\tilde{\eta}_{1q}^{L,R}$	0	–	✓
\tilde{S}_8 (spin-0 FV octet)	$\tilde{\eta}_{8q}^{L,R}$	0	+	×
S_3^α (spin-0 FV triplet)	η_3	–	0	×
$S_6^{\alpha\beta}$ (spin-0 FV sextet)	η_6	+	0	✓

$1-\sigma$ favored region for V_8



$$r_q = g_{8q}^R / g_{8q}^L \text{ and } g_{8q}^L = 1$$

Constraints on masses and couplings

- 1- σ favored values of the couplings **Updated data:**

$$\tilde{V}_8 : \frac{1}{N_c} \left(\frac{1 \text{ TeV}}{m_{\tilde{V}}} \right)^2 \left(|\tilde{g}_{8q}^L|^2 + |\tilde{g}_{8q}^R|^2 \right) \simeq 0.76(0.64),$$

$$\tilde{S}_1 : \left(\frac{1 \text{ TeV}}{m_{\tilde{S}}} \right)^2 \left(|\tilde{\eta}_{1q}^L|^2 + |\tilde{\eta}_{1q}^R|^2 \right) \simeq 0.62(0.49),$$

$$S_{13}^{\alpha\beta} : 2 \left(\frac{1 \text{ TeV}}{m_{S_6}} \right)^2 |\eta_6|^2 \simeq 0.76(0.64)$$

These could be discovered and tested at the LHC, by measuring the mass and the couplings

A_{FB} implies P violation

- If P were conserved in the light quark sector, $C_{8q}^{LL} = C_{8q}^{RL}$, and $C_{8q}^{LR} = C_{8q}^{RR}$
- If P were conserved in the top sector, $C_{8q}^{LL} = C_{8q}^{LR}$ and $C_{8q}^{RR} = C_{8q}^{RL}$
- In either case, we would have $C_1 - C_2 = 0$, and no effects on A_{FB}
- Most important message of nonzero A_{FB} from new physics is
Parity should be violated in the quark sector
- What would be the observable consequences of this new PV interaction ?
- In the top sector, the top longitudinal polarization can be nonzero in general, unlike QCD

$$\langle \vec{S}_t \cdot \hat{n} \rangle \neq 0$$

- If $\hat{n} \propto \vec{p}$, it becomes helicity

P violation and Longitudinal Pol's

- Keeping only the longitudinal pol's of (anti)top, we have

$$\overline{|\mathcal{M}|^2} = \frac{g_s^4}{\hat{s}^2} \left\{ \mathcal{D}_0 + \mathcal{D}_1(P_L + \bar{P}_L) + \mathcal{D}_2(P_L - \bar{P}_L) + \mathcal{D}_3 P_L \bar{P}_L \right\}.$$

- D_1 and D_2 are P-violating.
- D_1 term needs strong phase (CPT violating), from the propagator of the exchanged new particle, which is not seen in our approach (Future work)

•

$$\mathcal{D}_2 \simeq \frac{\hat{s}}{9\Lambda^2} \left[(C'_1 + C'_2) \hat{\beta}_t (1 + c_\theta^2) + (C'_1 - C'_2) (5 - 3\hat{\beta}_t^2) c_\theta \right]$$

with

$$C'_1 \equiv C_{8q}^{RR} - C_{8q}^{LL},$$

$$C'_2 \equiv C_{8q}^{LR} - C_{8q}^{RL}.$$

- Different informations on the chiral structures of NP from A_{FB}

- When we integrate over the polar angle $\hat{\theta}$, only the first term involving

$$(C'_1 + C'_2) = C_{8q}^{RR} - C_{8q}^{LL} + C_{8q}^{LR} - C_{8q}^{RL}$$

survives.

- On the other hand, if we separate the forward and the backward top samples and take the difference, the orthogonal combination in the second term survives:

$$(C'_1 - C'_2) = C_{8q}^{RR} - C_{8q}^{LL} - C_{8q}^{LR} + C_{8q}^{RL}.$$

- Consider the two new observables:

$$D \equiv \frac{\sigma(t_R \bar{t}_L) - \sigma(t_L \bar{t}_R)}{\sigma(t_R \bar{t}_R) + \sigma(t_L \bar{t}_L) + \sigma(t_L \bar{t}_R) + \sigma(t_R \bar{t}_L)},$$

$$D_{\text{FB}} \equiv D(\cos \hat{\theta} \geq 0) - D(\cos \hat{\theta} \leq 0)$$

which involve the sum and difference of the coefficients C'_1 and C'_2 , respectively.




Figure: The P -violating spin correlations D and D_{FB} in the (C'_1, C'_2) plane. The signs of (D, D_{FB}) are denoted.

- The polarization coefficients could be measured by studying the angular distributions of the top-quark decay products.
- For dilepton decay channels of $t\bar{t}$ (in the helicity basis), one has

$$|\overline{\mathcal{M}}|^2 = \frac{g_s^4}{\hat{s}^2} \left\{ \mathcal{D}_0 + \mathcal{D}_1(\cos \theta_+^* + \cos \theta_-^*) + \mathcal{D}_2(\cos \theta_+^* - \cos \theta_-^*) + \mathcal{D}_3 \cos \theta_+^* \cos \theta_-^* \right\}.$$

where θ_+^* (θ_-^*) is the angle between the charged lepton l^+ (l^-) in the top (anti-top) rest frame and the direction of the top (anti-top) in the $t\bar{t}$ rest frame.

- M_{T2} could be helpful for this study (Work in progress)

[t] *New particle exchanges and the signs of induced couplings C^{AB}*
($A, B = R, L$), $C_1 - C_2$, $C'_1 + C'_2$, and $C'_1 - C'_2$.

Res.	C^{RR}	C^{LL}	C^{LR}	C^{RL}	$C_1 - C_2$	$C'_1 + C'_2$	$C'_1 - C'_2$	
\tilde{V}_{1R}	-	0	0	0	-	-	-	
\tilde{V}_{1L}	0	-	0	0	-	+	+	
\tilde{V}_{8R}	+	0	0	0	+	+	+	
\tilde{V}_{8L}	0	+	0	0	+	-	-	
\tilde{S}_{1R}	0	0	0	-	+	+	-	
\tilde{S}_{1L}	0	0	-	0	+	-	+	
\tilde{S}_{8R}	0	0	0	+	-	-	+	
\tilde{S}_{8L}	0	0	+	0	-	+	-	
S_2^α	-	0	0	0	-	-	-	
$S_{13}^{\alpha\beta}$	+	0	0	0	+	+	+	
V_{8R}	\pm	0	0	0	\pm	\pm	\pm	




Figure: The predictions for D and D_{FB} of the models under consideration, being consistent with the $\sigma_{t\bar{t}}$ and A_{FB} measurements at the $1\text{-}\sigma$ level. We assume only one resonance exists or dominates.

Summary

- We performed a model independent study of $t\bar{t}$ productions at the Tevatron using dimension-6 $q\bar{q}t\bar{t}$ contact interactions with all the possible Dirac and color structures.
- Our results encode the necessary conditions for the underlying new physics in a compact and an effective way when those new particles are too heavy to be produced at the Tevatron.
- Proposed a new FB spin-spin correlation $C_{\text{FB}} \propto A_{\text{FB}}$
- Suggested to measure P_L and \overline{P}_L to probe the chiral structure of new physics in $q\bar{q} \rightarrow t\bar{t}$
- We considered the s -, t - and u -channel exchanges of spin-0 and spin-1 particles whose color quantum number is either singlet, octet, triplet or sextet.

Future Study

- A_{FB} vs $M_{t\bar{t}}$ or $\Delta\eta$
- Detailed study of resonance productions at the LHC
(Resonances in the $t\bar{t}$ or $t\bar{q}$ channel)
- Top polarization in general (transverse, longitudinal) at the Tevatron and at the LHC
- In the presence of resonance, there could be nonzero $P_L + \overline{P}_L \rightarrow$
Need more study the D_1 term
- Those new particles might leave imprints on the low energy flavor physics such as K or B physics (mixing and CP violation), if $u(d) - t$ transitions are employed

Thank you very much